York University

Faculties of Pure and Applied Science, Arts, Atkinson MATH 2090. Problem Set #1. Posted January 19, 2002

Due in the Tutorial of Friday, January 25, 2002

Section N

- In your proofs it is imperative to clearly state what **tools** you use (e.g., WLUS, SLCS, MP, PSL, Monotonicity, Deduction Theorem, Weak Generalization, which axiom(s), etc.)
 - **1.** (5 MARKS) In class we have proved that $\vdash (\exists x)(\forall y)A \Rightarrow (\forall y)(\exists x)A$ using the Auxiliary Variable Metatheorem.

Now prove the same thing but <u>without</u> the help of the Auxiliary Variable Metatheorem.

- **2.** (5 MARKS) Prove that $\vdash ((\forall x)B \Rightarrow A) \equiv (\exists x)(B \Rightarrow A)$, provided x is not free in A. Use an equational proof!
- **3.** (5 MARKS) Let f be a function symbol of arity 1. Prove that $\vdash x \approx y \Rightarrow fx \approx fy$.
- Careful! Do *not* just say that this "follows" from Chapter 1 stuff. It doesn't. This is a Predicate Calculus exercise!
 - 4. (5 MARKS) Prove the metatheorem "∀ commutativity (symmetry)" below.

$$\vdash (\forall x)(\forall y)A \equiv (\forall y)(\forall x)A$$

- 5. Let P be any predicate of arity 2.
 - (a) (2 MARKS) Explain WHY $(\forall x)(\forall y)P(x,y) \Rightarrow (\forall y)P(y,y)$ is NOT an instance of Ax2.
 - (b) (5 MARKS) Nevertheless, prove that $(\forall x)(\forall y)P(x,y) \Rightarrow (\forall y)P(y,y)$ IS an absolute theorem.
- **6.** (5 MARKS) Prove (absolutely) the formula $(\exists x)A \land (\forall x)B \Rightarrow (\exists x)(A \land B)$.
- 7. (5 MARKS) Prove (absolutely) the formula $(\exists x)(A \Rightarrow (\forall x)A)$
- \diamond A may have free x occurrences!
- 8. (5 MARKS) Let x, y, z be distinct variables and P, Q predicates of one variable. Prove the formula

$$Q(x) \Rightarrow (\forall y) \Big(Q(y) \Rightarrow (\forall z) P(z) \Big) \Rightarrow (\forall x) P(x)$$

Page 1

G. Tourlakis

ŚŚ

Ş

Ś

Ś

9. (5 MARKS) For any predicate P of arity 2 prove

$$\vdash (\forall x)(\forall y)P(x,y) \equiv (\forall y)(\forall x)P(y,x)$$

 \bigotimes This is different from problem #4. (How?)